

Non-symmetric convex domains have no basis of exponentials

MIHAIL N. KOLOUNTZAKIS^{1 2}

December 1998

Abstract

A conjecture of Fuglede states that a bounded measurable set $\Omega \subset \mathbb{R}^d$, of measure 1, can tile \mathbb{R}^d by translations if and only if the Hilbert space $L^2(\Omega)$ has an orthonormal basis consisting of exponentials $e_\lambda(x) = \exp 2\pi i \langle \lambda, x \rangle$. If Ω has the latter property it is called *spectral*. We generalize a result of Fuglede, that a triangle in the plane is not spectral, proving that every non-symmetric convex domain in \mathbb{R}^d is not spectral.

§0. Introduction

Let Ω be a measurable subset of \mathbb{R}^d of measure 1 and Λ be a discrete subset of \mathbb{R}^d . We write

$$\begin{aligned} e_\lambda(x) &= \exp 2\pi i \langle \lambda, x \rangle, \quad (x \in \mathbb{R}^d), \\ E_\Lambda &= \{e_\lambda : \lambda \in \Lambda\} \subset L^2(\Omega). \end{aligned}$$

The inner product and norm on $L^2(\Omega)$ are

$$\langle f, g \rangle_\Omega = \int_\Omega f \bar{g}, \quad \text{and} \quad \|f\|_\Omega^2 = \int_\Omega |f|^2.$$

Definition 1 The pair (Ω, Λ) is called a *spectral pair* if E_Λ is an orthonormal basis for $L^2(\Omega)$. A set Ω will be called *spectral* if there is $\Lambda \subset \mathbb{R}^d$ such that (Ω, Λ) is a spectral pair. The set Λ is then called a *spectrum* of Ω .

Example: If $Q_d = (-1/2, 1/2)^d$ is the cube of unit volume in \mathbb{R}^d then (Q_d, \mathbb{Z}^d) is a spectral pair.

We write $B_R(x) = \{y \in \mathbb{R}^d : |x - y| < R\}$.

Definition 2 (Density)

- (i) The set $\Lambda \subset \mathbb{R}^d$ has *uniformly bounded density* if for each $R > 0$ there exists a constant $C > 0$ such that Λ has at most C elements in each ball of radius R in \mathbb{R}^d .
- (ii) The set $\Lambda \subset \mathbb{R}^d$ has *density* ρ , and we write $\rho = \text{dens } \Lambda$, if for each $x \in \mathbb{R}^d$

$$\rho = \lim_{R \rightarrow \infty} \frac{|\Lambda \cap B_R(x)|}{|B_R|}.$$

We define translational tiling for complex-valued functions below.

Definition 3 Let $f : \mathbb{R}^d \rightarrow \mathbb{C}$ be measurable and $\Lambda \subset \mathbb{R}^d$ be a discrete set. We say that f *tiles with Λ at level $w \in \mathbb{C}$* , and sometimes write “ $f + \Lambda = w\mathbb{R}^d$ ”, if

$$\sum_{\lambda \in \Lambda} f(x - \lambda) = w, \quad \text{for almost every (Lebesgue) } x \in \mathbb{R}^d, \quad (1)$$

¹Department of Mathematics, 1409 W. Green St, University of Illinois, Urbana, IL 61801, USA. E-mail: kolount@math.uiuc.edu

²Partially supported by the U.S. National Science Foundation, under grant DMS 97-05775.

with the sum above converging absolutely a.e. If $\Omega \subset \mathbb{R}^d$ is measurable we say that $\Omega + \Lambda$ is a tiling when $\mathbf{1}_\Omega + \Lambda = w\mathbb{R}^d$, for some w . If w is not mentioned it is understood to be equal to 1.

Remarks

1. If $f \in L^1(\mathbb{R}^d)$ and Λ has uniformly bounded density one can easily show (see [KL96] for the proof in one dimension, which works in higher dimension as well) that the sum in (1) converges absolutely a.e. and defines a locally integrable function of x .
2. In the very common case when $f \in L^1(\mathbb{R}^d)$ and $\int_{\mathbb{R}^d} f \neq 0$ the condition that Λ has uniformly bounded density follows easily from (1) and need not be postulated a priori.
3. It is easy to see that if $f \in L^1(\mathbb{R}^d)$, $\int_{\mathbb{R}^d} f \neq 0$ and $f + \Lambda$ is a tiling then Λ has a density and the level of the tiling w is given by

$$w = \int_{\mathbb{R}^d} f \cdot \text{dens } \Lambda.$$

From now on we restrict ourselves to tiling with functions in L^1 and sets of finite measure.

Example: $Q_d + \mathbb{Z}^d$ is a tiling.

The following conjecture is still unresolved.

Conjecture: (Fuglede [F74]) If $\Omega \subset \mathbb{R}^d$ is bounded and has Lebesgue measure 1 then $L^2(\Omega)$ has an orthonormal basis of exponentials if and only if there exists $\Lambda \subset \mathbb{R}^d$ such that $\Omega + \Lambda = \mathbb{R}^d$ is a tiling.

It is not hard to show [F74] that $L^2(\Omega)$ has a basis Λ which is a *lattice* (i.e., $\Lambda = A\mathbb{Z}^d$, where A is a non-singular $d \times d$ matrix) if and only if $\Omega + \Lambda^*$ is a tiling. Here

$$\Lambda^* = \left\{ \mu \in \mathbb{R}^d : \langle \mu, \lambda \rangle \in \mathbb{Z}, \forall \lambda \in \Lambda \right\}$$

is the *dual lattice* of Λ (we have $\Lambda^* = A^{-\top}\mathbb{Z}^d$).

Fuglede [F74] showed that the disk and the triangle in \mathbb{R}^2 are not spectral domains.

In this note we prove the following generalization of Fuglede’s triangle result.

Theorem 1 *Let Ω have measure 1 and be a convex, non-symmetric, bounded open set in \mathbb{R}^d . Then Ω is not spectral.*

The set Ω is called *symmetric* with respect to 0 if $y \in \Omega$ implies $-y \in \Omega$, and symmetric with respect to $x_0 \in \mathbb{R}^d$ if $y \in \Omega$ implies that $2x_0 - y \in \Omega$. It is called *non-symmetric* if it is not symmetric with respect to any $x_0 \in \mathbb{R}^d$. For example, in any dimension a simplex is non-symmetric.

It is known [V54, M80] that every convex body that tiles \mathbb{R}^d by translation is a centrally symmetric polytope and that each such body also admits a lattice tiling and, therefore (see the remark after Fuglede’s conjecture above), its L^2 admits a lattice spectrum. Given Theorem 1, to prove Fuglede’s conjecture restricted to convex domains, one still has to prove that any symmetric convex body that is not a tile admits no orthonormal basis of exponentials for its L^2 .

In §1 we derive some necessary and some sufficient conditions for $f + \Lambda$ to be a tiling. These conditions roughly state that tiling is equivalent to a certain tempered distribution, associated with Λ , being “supported” on the zero set of \widehat{f} plus the origin. Similar conditions had been derived in [KL96] but here we have to work with less smoothness for \widehat{f} . To

compensate for the lack of smoothness we work with compactly supported \widehat{f} and nonnegative f and \widehat{f} , conditions which are fulfilled for our problem.

In §2 we restate the property that Ω is spectral as a tiling problem for $|\widehat{\mathbf{1}}_\Omega|^2$ and use the conditions derived in §1 to prove Theorem 1. What makes the proof work is that, when Ω is a non-symmetric convex set, the set $\Omega - \Omega$ has volume strictly larger than $2^d \text{vol } \Omega$.

§1. Fourier-analytic conditions for tiling

Our method relies on a Fourier-analytic characterization of translational tiling, which is a variation of the one used in [KL96]. We define the (generally unbounded) measure

$$\delta_\Lambda = \sum_{\lambda \in \Lambda} \delta_\lambda,$$

where δ_λ represents a unit mass at $\lambda \in \mathbb{R}^d$. If Λ has uniformly bounded density then $\widehat{\delta}_\Lambda$ is a tempered distribution (see for example [R73]) and therefore its Fourier Transform $\widehat{\delta}_\Lambda$ is defined and is itself a tempered distribution.

The action of a tempered distribution α on a Schwartz function ϕ is denoted by $\langle \phi, \alpha \rangle$. The Fourier Transform of α is defined by the equation

$$\langle \phi, \widehat{\alpha} \rangle = \langle \widehat{\phi}, \alpha \rangle.$$

The support $\text{supp } \alpha$ is the smallest closed set F such that for any smooth ϕ of compact support contained in the open set F^c we have $\langle \phi, \alpha \rangle = 0$.

Theorem 2 *Suppose that $f \geq 0$ is not identically 0, that $f \in L^1(\mathbb{R}^d)$, $\widehat{f} \geq 0$ has compact support and $\Lambda \subset \mathbb{R}^d$. If $f + \Lambda$ is a tiling then*

$$\text{supp } \widehat{\delta}_\Lambda \subseteq \{x \in \mathbb{R}^d : \widehat{f}(x) = 0\} \cup \{0\}. \quad (2)$$

Proof. Assume that $f + \Lambda = w\mathbb{R}^d$ and let

$$K = \{\widehat{f} = 0\} \cup \{0\}.$$

We have to show that

$$\langle \phi, \widehat{\delta}_\Lambda \rangle = 0, \quad \forall \phi \in C_c^\infty(K^c).$$

This is equivalent to $\sum_{\lambda \in \Lambda} \widehat{\phi}(\lambda) = 0$, for each such ϕ . Notice that $h = \phi/\widehat{f}$ is a continuous function, however not necessarily smooth. We shall need that $\widehat{h} \in L^1$. This is a consequence of Wiener's Tauberian Theorem [R73, Ch. 11]. We denote by $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$ the d -dimensional torus.

(Wiener) *If $g \in C(\mathbb{T}^d)$ has an absolutely convergent Fourier series*

$$g(x) = \sum_{n \in \mathbb{Z}^d} \widehat{g}(n) e^{2\pi i \langle n, x \rangle}, \quad \widehat{g} \in \ell^1(\mathbb{Z}^d),$$

and if g does not vanish anywhere on \mathbb{T}^d then $1/g$ also has an absolutely convergent Fourier series.

Assume that

$$\text{supp } \widehat{f} \subseteq \left(-\frac{L}{2}, \frac{L}{2}\right)^d.$$

Define the function F to be:

- (i) periodic in \mathbb{R}^d with period lattice $(L\mathbb{Z})^d$,
- (ii) to agree with \widehat{f} on $\text{supp } \phi$,
- (iii) to be non-zero everywhere and,
- (iv) to have $\widehat{F} \in \ell^1(\mathbb{Z}^d)$, i.e.,

$$\widehat{F} = \sum_{n \in \mathbb{Z}^d} \widehat{F}(n) \delta_{L^{-1}n},$$

is a finite measure in \mathbb{R}^d .

One way to define such an F is as follows. First, define the $(L\mathbb{Z})^d$ -periodic function $g \geq 0$ to be \widehat{f} periodically extended. The Fourier coefficients of g are $\widehat{g}(n) = L^{-d} f(-n/L) \geq 0$. By the Poisson Summation Formula

$$\begin{aligned} \sum_{n \in \mathbb{Z}^d} |\widehat{g}(n)| &= \sum_{n \in \mathbb{Z}^d} L^{-d} f\left(\frac{n}{L}\right) \\ &= \sum_{n \in \mathbb{Z}^d} \widehat{f}(Ln) < \infty \end{aligned}$$

since \widehat{f} has compact support.

Let ϵ be small enough to guarantee that \widehat{f} (and hence g) does not vanish on $\text{supp } \phi + B_\epsilon(0)$. Let k be a smooth $(L\mathbb{Z})^d$ -periodic function which is equal to 1 on $\text{supp } \phi$ and equal to 0 off $\text{supp } \phi + B_\epsilon(0)$, and satisfies $0 \leq k \leq 1$ everywhere. Finally, define

$$F = kg + (1 - k).$$

Since both k and g have absolutely summable Fourier series and this property is preserved under both sums and products, it follows that F also has an absolutely summable Fourier series. And by the nonnegativity of g we get that F is never 0.

By Wiener's theorem, $\widehat{F^{-1}} \in \ell^1(\mathbb{Z}^d)$, i.e., $\widehat{F^{-1}}$ is a finite measure on \mathbb{R}^d . We now have that

$$\left(\frac{\phi}{\widehat{f}}\right)^\wedge = \widehat{\phi F^{-1}} = \widehat{\phi} * \widehat{F^{-1}} \in L^1(\mathbb{R}^d).$$

This justifies the interchange of the summation and integration below:

$$\begin{aligned} \sum_{\lambda \in \Lambda} \widehat{\phi}(\lambda) &= \sum_{\lambda \in \Lambda} \int \frac{\phi(x)}{\widehat{f}(x)} \widehat{f}(x) e^{-2\pi i \langle \lambda, x \rangle} dx \\ &= \sum_{\lambda \in \Lambda} \left(\frac{\phi}{\widehat{f}}\right)^\wedge(\lambda) \\ &= \sum_{\lambda \in \Lambda} \left(\frac{\phi}{\widehat{f}}\right)^\wedge * \widehat{f}(\lambda) \\ &= \sum_{\lambda \in \Lambda} \int_{\mathbb{R}^d} \left(\frac{\phi}{\widehat{f}}\right)^\wedge(y) f(y - \lambda) dy \\ &= \int_{\mathbb{R}^d} \left(\frac{\phi}{\widehat{f}}\right)^\wedge(y) \sum_{\lambda \in \Lambda} f(y - \lambda) dy \\ &= w \int_{\mathbb{R}^d} \left(\frac{\phi}{\widehat{f}}\right)^\wedge(y) \\ &= 0, \end{aligned}$$

as we had to show.

■

For a set $A \subseteq \mathbb{R}^d$ and $\delta > 0$ we write

$$A_\delta = \left\{ x \in \mathbb{R}^d : \text{dist}(x, A) < \delta \right\}.$$

We shall need the following partial converse to Theorem 2.

Theorem 3 *Suppose that $f \in L^1(\mathbb{R}^d)$, and that $\Lambda \subset \mathbb{R}^d$ has uniformly bounded density. Suppose also that $O \subset \mathbb{R}^d$ is open and*

$$\text{supp } \widehat{\delta_\Lambda} \setminus \{0\} \subseteq O \text{ and } O_\delta \subseteq \left\{ \widehat{f} = 0 \right\}, \quad (3)$$

for some $\delta > 0$. Then $f + \Lambda$ is a tiling at level $\widehat{f}(0) \cdot \widehat{\delta_\Lambda}(\{0\})$.

Proof. Let $\psi : \mathbb{R}^d \rightarrow \mathbb{R}$ be smooth, have support in $B_1(0)$ and $\widehat{\psi}(0) = 1$ and for $\epsilon > 0$ define the approximate identity $\psi_\epsilon(x) = \epsilon^{-d}\psi(x/\epsilon)$. Let

$$f_\epsilon = \widehat{\psi}_\epsilon f,$$

which has rapid decay.

First we show that $(\int f_\epsilon)^{-1} f_\epsilon + \Lambda$ is a tiling. That is, we show that the convolution $f_\epsilon * \delta_\Lambda$ is a constant. Let ϕ be any Schwartz function. Then

$$\langle \phi, f_\epsilon * \delta_\Lambda \rangle = \langle \widehat{\phi}, \widehat{f_\epsilon} \widehat{\delta_\Lambda} \rangle = \langle \widehat{\phi} \widehat{f_\epsilon}, \widehat{\delta_\Lambda} \rangle.$$

The function $\widehat{\phi} \widehat{f_\epsilon}$ is a Schwartz function whose support intersects $\text{supp } \widehat{\delta_\Lambda}$ only at 0, since, for small enough $\epsilon > 0$,

$$\text{supp } \widehat{\phi} \widehat{f_\epsilon} \subseteq \text{supp } \widehat{f_\epsilon} \subseteq (\text{supp } \widehat{f})_\epsilon \subseteq O^c.$$

Hence, for each Schwartz function ϕ

$$\langle \phi, f_\epsilon * \delta_\Lambda \rangle = \widehat{\phi}(0) \widehat{f_\epsilon}(0) \widehat{\delta_\Lambda}(\{0\}),$$

which implies

$$f_\epsilon * \delta_\Lambda(x) = \widehat{f_\epsilon}(0) \widehat{\delta_\Lambda}(\{0\}), \text{ a.e.}(x).$$

We also have that $\sum_{\lambda \in \Lambda} |f(x - \lambda)|$ is finite a.e. (see Remark 1 following the definition of tiling), hence, for almost every $x \in \mathbb{R}^d$

$$\sum_{\lambda \in \Lambda} |f(x - \lambda) - f_\epsilon(x - \lambda)| = \sum_{\lambda \in \Lambda} |f(x - \lambda)| \cdot \left| 1 - \widehat{\psi}_\epsilon(x - \lambda) \right|,$$

which tends to 0 as $\epsilon \rightarrow 0$. This proves

$$\sum_{\lambda \in \Lambda} f(x - \lambda) = \widehat{f}(0) \cdot \widehat{\delta_\Lambda}(\{0\}), \text{ a.e.}(x).$$

■

§2. Proof of the main result

We now make some remarks that relate the property of E_Λ being a basis for $L^2(\Omega)$ to a certain function tiling \mathbb{R}^d with Λ .

Assume that Ω is a bounded set of measure 1. Notice first that

$$\langle e_\lambda, e_x \rangle_\Omega = \widehat{\mathbf{1}}_\Omega(x - \lambda).$$

The set E_Λ is an orthonormal basis for $L^2(\Omega)$ if and only if for each $f \in L^2(\Omega)$

$$\|f\|_\Omega^2 = \sum_{\lambda \in \Lambda} |\langle e_\lambda, f \rangle_\Omega|^2,$$

and, by the completeness of the exponentials in L^2 of a large cube containing Ω , it is necessary and sufficient that

$$\sum_{\lambda \in \Lambda} \left| \widehat{\mathbf{1}}_\Omega(x - \lambda) \right|^2 = 1, \quad (4)$$

for each $x \in \mathbb{R}^d$. In other words a necessary and sufficient condition for (Ω, Λ) to be a spectral pair is that $\left| \widehat{\mathbf{1}}_\Omega \right|^2 + \Lambda$ is a tiling at level 1. Notice also that $\left| \widehat{\mathbf{1}}_\Omega \right|^2$ is the Fourier Transform of $\mathbf{1}_\Omega * \widetilde{\mathbf{1}}_\Omega$ which has support equal to the set $\Omega - \Omega$. We use the notation $\widehat{f}(x) = \overline{f(-x)}$.

Proof of Theorem 1: Write $K = \Omega - \Omega$, which is a symmetric convex set. Assume that (Ω, Λ) is a spectral pair. We can clearly assume that $0 \in \Lambda$. It follows that $\left| \widehat{\mathbf{1}}_\Omega \right|^2 + \Lambda$ is a tiling and hence that Λ has uniformly bounded density, has density equal to 1 and $\widehat{\delta}_\Lambda(\{0\}) = 1$.

By Theorem 2 with

$$f = \left| \widehat{\mathbf{1}}_\Omega \right|^2, \quad \widehat{f} = \mathbf{1}_\Omega * \widetilde{\mathbf{1}}_\Omega(-x)$$

it follows that

$$\text{supp } \widehat{\delta}_\Lambda \subseteq \{0\} \cup K^c.$$

Let $H = K/2$ and write

$$f(x) = \mathbf{1}_H * \widetilde{\mathbf{1}}_H(x) = \int_{\mathbb{R}^d} \mathbf{1}_H(y) \mathbf{1}_H(y - x) dy.$$

The function f is supported in K and has nonnegative Fourier Transform

$$\widehat{f} = \left| \widehat{\mathbf{1}}_H \right|^2.$$

We have

$$\int_{\mathbb{R}^d} \widehat{f} = f(0) = \text{vol } H$$

and

$$\widehat{f}(0) = \int_{\mathbb{R}^d} f = (\text{vol } H)^2.$$

By the Brunn-Minkowski inequality (see for example [G94, Ch. 3]), for any convex body Ω ,

$$\text{vol } \frac{1}{2}(\Omega - \Omega) \geq \text{vol } \Omega,$$

with equality only in the case of symmetric Ω . Since Ω has been assumed to be non-symmetric it follows that

$$\text{vol } H > 1.$$

For

$$1 > \rho > \left(\frac{1}{\text{vol } H} \right)^{1/d}$$

consider

$$g(x) = f(x/\rho)$$

which is supported properly inside K , and has

$$g(0) = f(0) = \text{vol } H, \quad \int_{\mathbb{R}^d} g = \rho^d \int_{\mathbb{R}^d} f = \rho^d (\text{vol } H)^2.$$

Since $\text{supp } g$ is properly contained in K Theorem 3 implies that $\widehat{g} + \Lambda$ is a tiling at level $\int \widehat{g} \cdot \text{dens } \Lambda = \int \widehat{g} = g(0) = \text{vol } H$. However, the value of \widehat{g} at 0 is $\int g = \rho^d (\text{vol } H)^2 > \text{vol } H$, and, since $\widehat{g} \geq 0$ and \widehat{g} is continuous, this is a contradiction.

■

§3. Bibliography

- [F74] B. Fuglede, Commuting self-adjoint partial differential operators and a group theoretic problem, *J. Funct. Anal.* **16**(1974), 101-121.
- [G94] R.J. Gardner, *Geometric tomography*, Cambridge University Press, 1994.
- [KL96] M.N. Kolountzakis and J.C. Lagarias, Structure of tilings of the line by a function, *Duke Math. J.* **82** (1996), 3, 653-678.
- [M80] P. McMullen, Convex bodies which tile space by translation, *Mathematika* **27** (1980), 113-121.
- [R73] W. Rudin, *Functional analysis*, McGraw-Hill, New York, 1973.
- [V54] B.A. Venkov, On a class of Euclidean polyhedra, *Vestnik Leningrad Univ. Ser. Math. Fiz. Him.* **9** (1954), 11-31 (Russian).